

§ 180.421

[51 FR 12146, Apr. 9, 1986, as amended at 55 FR 29829, July 20, 1990]

§ 180.421 Fenarimol; tolerances for residues.

(a) General. (1) Tolerances are established for residues of the fungicide fenarimol [alpha-(2-chlorophenyl)-alpha-(4-chlorophenyl)-5-pyrimidinemethanol] in or on the following raw agricultural commodities:

Commodity	Parts per million
Apple	0.1
Apple, dry pomace	2.0
Apple, wet pomace	2.0
Cattle, fat	0.1
Cattle, kidney	0.1
Cattle, meat	0.01
Cattle, meat byproducts, except kidney	0.05
Goat, fat	0.1
Goat, kidney	0.1
Goat, meat	0.01
Goat, meat byproducts, except kidney	0.05
Horse, fat	0.1
Horse, kidney	0.1
Horse, meat	0.01
Horse, meat byproducts, except kidney	0.05
Pear	0.1
Pecan	0.1
Sheep, fat	0.1
Sheep, kidney	0.1
Sheep, meat	0.01
Sheep, meat byproducts, except kidney	0.05

(2) A tolerance is established for combined residues of the fungicide fenarimol [alpha-(2-chlorophenyl)-alpha-(4-chlorophenyl)-5-pyrimidinemethanol] and its metabolites [alpha-(2-chlorophenyl)-alpha-(4-chlorophenyl)-1,4-dihydro-5-pyrimidinemethanol and 5-[(2-chlorophenyl)(4-chlorophenyl)methyl]-3,4-dihydro-4-pyrimidinol measured as the total of fenarimol and 5-[(2-chlorophenyl)-(4-chlorophenyl)methyl]pyrimidine (calculated as fenarimol)], in or on the following raw agricultural commodities:

Commodity	Parts per million
Banana ¹	0.5
(Not more than 0.25 ppm shall be present in the pulp after peel is removed)	
Cherry	1.0
Grape, juice	0.6
Grape pomace (wet and dry)	2.0
Grape	0.2
Grape, raisin, waste	3.0
Grape, raisin	0.6

¹There are no United States registrations for banana as of April 26, 1995.

40 CFR Ch. I (7-1-04 Edition)

(b) Section 18 emergency exemptions. [Reserved]

(c) Tolerances with regional registrations. [Reserved]

(d) Indirect or inadvertent residues. [Reserved]

[51 FR 39662, Oct. 30, 1986, as amended at 53 FR 27349, July 20, 1988; 53 FR 44403, Nov. 3, 1988; 54 FR 45734, Oct. 31, 1989; 60 FR 33354, June 28, 1995; 62 FR 49937, Sept. 24, 1997; 62 FR 61447, Nov. 18, 1997; 67 FR 35048, May 17, 2002; 67 FR 41807, June 19, 2002; 69 FR 6567, Feb. 11, 2004]

§ 180.422 Tralomethrin; tolerances for residues.

(a) General. (1) Tolerances are established for the combined residues of the pesticide chemical tralomethrin ((S)-alpha-cyano-3-phenoxybenzyl (1R,3S)-2,2-dimethyl-3-[(RS)-1,2,2,2-tetrabromoethyl]-cyclopropanecarboxylate) and its metabolites (S)-alpha-cyano-3-phenoxybenzyl (1R,3R)-3-(2,2-dibromovinyl)-2,2-dimethyl-cyclopropanecarboxylate and (S)-alpha-cyano-3-phenoxybenzyl(1S,3R)-3-(2,2-dibromovinyl)-2,2-dimethyl-cyclopropanecarboxylate calculated as the parent in or on the following food commodities:

Commodity	Parts per million
Broccoli	0.5
Cotton, undelinted seed	0.02
Cotton, oil	0.20
Lettuce, head	1.00
Lettuce, leaf	3.00
Soybean	0.05
Sunflower, seed	0.05

(2) A tolerance of 0.02 part per million is established for the combined residues of the insecticide tralomethrin ((S)-alpha-cyano-3-phenoxybenzyl-(1R,3S)-2,2-dimethyl-3-[(RS)-1,2,2,2-tetrabromoethyl]-cyclopropanecarboxylate) and its metabolites cis-deltamethrin [(S)-alpha-cyano-3-phenoxybenzyl-(1R,3R)-3-(2,2-dibromovinyl)-2,2-dimethylcyclopropanecarboxylate] and trans-deltamethrin [(S)-alpha-cyano-3-phenoxybenzyl (1S,3R)-3-(2,2-dibromovinyl)-2,2-dimethylcyclopropanecarboxylate] as follows:

(i) In or on all food items (other than those covered by a higher tolerance as

Environmental Protection Agency

§ 180.423

a result of use on growing crops) in food-handling establishments.

(ii) The insecticide may be present as a residue from application of tralomethrin in food-handling establishments, including food service, manufacturing, and processing establishments, such as restaurants, cafeterias, supermarkets, bakeries, breweries, dairies, meat slaughtering and packing plants, and canneries in accordance with the following prescribed conditions:

(A) Application shall be limited to a general surface and spot and/or crack and crevice treatment in food-handling establishments where food and food products are held, processed, prepared, and served. General surface application may be used only when the facility is not in operation provided exposed food has been covered or removed from the area being treated. All food-contact surfaces and equipment must be thoroughly cleaned after general surface applications. Spot and/or crack and crevice application may be used while the facility is in operation provided exposed food is covered or removed from the area being treated prior to application. Spray concentration shall be limited to a maximum of 0.06 percent active ingredient. Contamination of food and food-contact surfaces shall be avoided.

(B) To assure safe use of the insecticide, its label and labelling shall conform to that registered with the U.S. Environmental Protection Agency and shall be used in accordance with such label and labelling.

(3) A tolerance of 0.02 part per million is established for the combined residues of the insecticide tralomethrin ((*S*)-*alpha*-cyano-3-phenoxybenzyl-(1*R*,3*S*)-2,2-dimethyl-3-[(*R**S*)-1,2,2,2-tetrabromoethyl]cyclopropanecarboxylate) and its metabolites *cis*-deltamethrin [(*S*)-*alpha*-cyano-3-phenoxybenzyl-(1*R*,3*R*)-3-(2,2-dibromovinyl)-2,2-dimethylcyclopropanecarboxylate] and *trans*-deltamethrin [(*S*)-*alpha*-cyano-3-phenoxybenzyl (1*S*,3*R*)-3-(2,2-dibromovinyl)-2,2-dimethylcyclopropanecarboxylate] as follows:

(i) In or on all feed items (other than those covered by a higher tolerance as

a result of use on growing crops) in feed-handling establishments.

(ii) The insecticide may be present as a residue from application of tralomethrin in feed-handling establishments, including feed manufacturing and processing establishments in accordance with the following prescribed conditions:

(A) Application shall be limited to a general surface and spot and/or crack and crevice treatment in feed-handling establishments where feed and feed products are held or processed. General surface application may be used only when the facility is not in operation provided exposed feed has been covered or removed from the area being treated. All feed-contact surfaces and equipment must be thoroughly cleaned after general surface applications. Spot and/or crack and crevice application may be used while the facility is in operation provided exposed feed is covered or removed from the area being treated prior to application. Spray concentration shall be limited to a maximum of 0.06 percent active ingredient. Contamination of feed and feed-contact surfaces shall be avoided.

(B) To assure safe use of the insecticide, its label and labelling shall conform to that registered with the U.S. Environmental Protection Agency and shall be used in accordance with such label and labelling.

(b) *Section 18 emergency exemptions.* [Reserved]

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.* [Reserved]

[62 FR 63001, Nov. 26, 1997, as amended at 62 FR 66025, Dec. 17, 1997; 65 FR 33701, May 24, 2000]

§ 180.423 Fenridazon, potassium salt; tolerances for residues.

Tolerances are established for residues of the hybridizing agent potassium salt of fenridazon (1-(4-chlorophenyl)-1,4-dihydro-6-methyl-4-oxo-3-pyridazinecarboxylic acid, potassium salt; CAS Reg. No. 83588-43-6) in or on the following raw agricultural commodities: